


Definition: Let $A \subset \mathbb{R}$ be a set. We say that


- (i) A is open if $\forall x \in A, \exists \delta > 0: \underline{(x-\delta, x+\delta) \subset A}$ or $A = \emptyset$
- (ii) A is closed if $\mathbb{R} \setminus A$ is open
- (iii) A is bounded if $\exists a, b \in \mathbb{R}, a < b: A \subset (a, b)$.
- (iv) A is unbounded if it is not bounded.

Examples:

the interval $(a, b) = \{x \in \mathbb{R}: a < x < b\}$ is open.




the set $(0, 1) \cup (2, 3)$ is open.



$\mathbb{R}$ is open: $x \in \mathbb{R}, \delta > 0 \Rightarrow (x-\delta, x+\delta) \subset \mathbb{R}$.

$\emptyset$ is open.

$A = \{x \in \mathbb{R}: 0 < x \leq 1\}$




Is A open? No ($1 \in A$)

$\mathbb{R}$ is closed (because $\mathbb{R} \setminus \mathbb{R} = \emptyset$ is open).

$\emptyset$ is closed (because $\mathbb{R}$ is open).


$A = \{x \in \mathbb{R}: a \leq x \leq b\}, a < b$, is closed



$$\mathbb{R} \setminus A = \{x \in \mathbb{R} : x < a \vee x > b\} = \underbrace{(-\infty, a) \cup (b, +\infty)}_{\text{this set is open.}}$$

• Any finite set is closed.

$A = \{x_1, \dots, x_n\}$ is closed




$$n = 3$$

$$\begin{aligned} \mathbb{R} \setminus A &= \{x \in \mathbb{R} : x \notin A\} = \{x \in \mathbb{R} : x \neq x_1, x \neq x_2, x \neq x_3\} \\ &= \underbrace{(-\infty, x_1)}_{\text{open}} \cup \underbrace{(x_1, x_2)}_{\text{open}} \cup \underbrace{(x_2, x_3)}_{\text{open}} \cup \underbrace{(x_3, +\infty)}_{\text{open}} = \text{open} \end{aligned}$$

general property: the union of open sets is open

• the interval $[a, +\infty) = \{x \in \mathbb{R} : x \geq a\}$ is closed.

$$\mathbb{R} \setminus [a, +\infty) = (-\infty, a) \text{ is open}$$



• the interval $(-\infty, a] = \{x \in \mathbb{R} : x \leq a\}$ is closed.


$$\mathbb{R} \setminus (-\infty, a] = (a, +\infty) \text{ is open.}$$

Boundedness:

• the interval (a, b) is bounded.

• the interval $[a, b]$ is bounded

$$[a, b] \subset (a-1, b+1)$$



... interval $(a-1, b+1) = \{x \in \mathbb{R} : a-1 < x < b+1\}$ and


$$a-1 \sim a \sim b \sim b+1$$

the intervals $[a, b) := \{x \in \mathbb{R} : a \leq x < b\}$ and $[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\}$ are bounded.

Finite sets are bounded.

$$A = \{x_1, \dots, x_n\} \subset (x_1 - 1, x_n + 1)$$


$$\text{assume } x_1 < \dots < x_i < \dots < x_n$$



A finite union of bounded set is bounded.

$$\bigcup_{n \in \mathbb{Z}} (n, n+1) = \mathbb{R} \setminus \mathbb{Z} \text{ is unbounded}$$

bounded $\forall n \in \mathbb{Z}$



the interval $(a, +\infty) = \{x \in \mathbb{R} : x > a\}$ is unbounded

the interval $[a, +\infty)$ is unbounded.

the intervals $(-\infty, a)$ and $(-\infty, a]$ are unbounded.

$\mathbb{R} = (-\infty, \infty)$ is unbounded, $\emptyset$ is bounded.

$$\mathbb{R} \not\subset (a, b) \quad \forall a, b \in \mathbb{R}.$$

Classification of the intervals:

(i) (a, b) , $a, b \in \mathbb{R}$: open bounded intervals.


- (i) (a, b) , $a, b \in \mathbb{R}$: open bounded intervals.
- (ii) $[a, b)$, $a, b \in \mathbb{R}$: bounded intervals, neither closed nor open.
- (iii) $[a, b]$, $a, b \in \mathbb{R}$: bounded intervals, neither closed nor open.
- (iv) $[a, b]$, $a, b \in \mathbb{R}$: closed bounded intervals.
- (v) (a, ∞) , $a \in \mathbb{R}$: open unbounded intervals.
- (vi) $[a, \infty)$, $a \in \mathbb{R}$: closed unbounded intervals.
- (vii) $(-\infty, a)$, $a \in \mathbb{R}$: open unbounded intervals.
- (viii) $(-\infty, a]$, $a \in \mathbb{R}$: open unbounded intervals.
- (ix) $\mathbb{R} = (-\infty, \infty)$ open and closed unbounded interval

Definition: Let $A \subset \mathbb{R}$ be a set. We say that a point $x \in A$ is an interior point if $\exists \delta > 0: (x - \delta, x + \delta) \subset A$.

$\overset{\circ}{A} := \{x \in A: x \text{ is an interior point of } A\}$.

($\overset{\circ}{A}$ is known as the interior of A).

$$A = (a, b]; \quad \overset{\circ}{A} = (a, b)$$



property: $\overset{\circ}{A} \subset A$.

The extrema of a set

Let $A \subset \mathbb{R}$ be a set.

Definition: we say that a point $x \in A$ is the maximum of A if $x \geq y, \forall y \in A$, and we write $x = \max A$.

• we say that a point $x \in A$ is the minimum of A if $x \leq y, \forall y \in A$, and we write $x = \min A$.

examples: $A = [0, 1] \Rightarrow 1 = \max A$

- $1 \in A$
- $1 \geq y, \forall y \in A$.

$$0 = \min A$$

- $0 \in A$
- $0 \leq y, \forall y \in A$

in particular, $[0, 1]$ has both a min and a max.

• $A = [0, 1] \quad \cdot \quad 1 = \max A. \quad (1 \in A, 1 \geq y, \forall y \in A).$

• $0 \leq y, \forall y \in A; \quad 0 \in A? \text{ No}$

0 is not the min of A .

• A has no minimum.

take $x \in A \Rightarrow x > 0; \quad \frac{x}{2} > 0 \Rightarrow \frac{x}{2} \in A$

$\frac{x}{2} < x \Rightarrow x$ is not the min of A .

• there are sets $A \subset \mathbb{R}$ which have no

maximum. $A = [0, 1)$

• there are sets $A \subset \mathbb{R}$ which have neither a min nor a max. For example $(0, 1)$ or $(0, \infty)$

• Given a set $A \subset \mathbb{R}$, we define the set

$$M(A) := \{x \in \mathbb{R} : x \geq y, \forall y \in A\}$$

and

$$m(A) := \{x \in \mathbb{R} : x \leq y, \forall y \in A\}.$$

The set $M(A)$ is known as the set of majorants of A , while the set $m(A)$ is known as the set of minorants of A .

$$A = [0, 1] \quad \text{---} \frac{0 \quad 1}{[\quad]} \text{---}$$

$$M(A) = [1, \infty), \quad m(A) = (-\infty, 0]$$

$$1 \geq x, \forall x \in A$$

$$0 \leq x, \forall x \in A.$$

$$A = (0, \infty) \Rightarrow M(A) = \emptyset.$$

Theorem Let $A \subset \mathbb{R}$ be a set.

Then $M(A)$ is empty or it has a min.

and $m(A)$ is empty or it has a max.

equivalently: if $M(A) \neq \emptyset \Rightarrow M(A)$ has a min

$\neg \exists m(A \neq \emptyset \Rightarrow m(A) \text{ has a max.})$

As a consequence, we are allowed to introduce the following def:

Definition: let $A \subset \mathbb{R}$ be a set.

$$\sup(A) := \begin{cases} +\infty & \text{if } M(A) = \emptyset \\ \min M(A) & \text{if } M(A) \neq \emptyset \end{cases}$$

$$\inf(A) := \begin{cases} -\infty & \text{if } m(A) = \emptyset \\ \max m(A) & \text{if } m(A) \neq \emptyset \end{cases}$$

In particular: any subset of $\mathbb{R}$ has both an inf and a sup.

$$A = (0, 1] , \quad 0 = \inf A.$$

$$m(A) = (-\infty, 0] \Rightarrow 0 = \max m(A).$$

$$A = [0, 1) \Rightarrow 1 = \sup A$$

$$\sup(a, b) = \sup[a, b] = \sup[a, b) = b$$

||

$$\sup[a, b).$$

Theorem: let $A \subset \mathbb{R}$ be a set. Then

(i) A has a max $\Leftrightarrow \sup A \in A$ and, in this case, $\sup A = \max A$.

(ii) A has a min $\Leftrightarrow \inf A \in A$ and, in this

case, $\sup A = \max A$.

(ii) A has a min ($\Rightarrow$) $\inf A \in A$ and, in this case, $\inf A = \min A$.

Remark: the sup of a set can be $+\infty$ while the max cannot.

the inf of a set can be $-\infty$, while the min cannot.

Examples:

$$\max (a, b] = \max [a, b] = b : \quad \begin{array}{l} b \in (a, b] \\ b \in [a, b] \end{array}$$

(a, b) has no max.

$$\sup (a, b) = b$$

$$\sup (0, \infty) = +\infty$$

$$\inf (-\infty, 0) \stackrel{?}{=} -\infty$$

$$\inf (a, b) = a$$

Roughly speaking, if the extrema of A belong to A , we call them max and min. If they do not belong to A , we call them inf and sup.

By definition: $A \subset \mathbb{R}$ is bounded if and only

By definition: $A \subset \mathbb{R}$ is bounded if and only if $A \subset (a, b)$ for some $a, b \in \mathbb{R}$.

Remark: $A \neq \emptyset$ is bounded $(\Leftrightarrow) -\infty < \inf A \leq \sup A < \infty$
 $(\Leftrightarrow) \inf A, \sup A \in \mathbb{R}$.

Definition: $\emptyset \neq A \subset \mathbb{R}$ is bounded below if $\inf A > -\infty$
is bounded above if $\sup A < +\infty$.

Example: $(0, \infty)$ is bounded below.
 $(-\infty, 1)$ is bounded above.

Remark: if A is bounded $(\Leftrightarrow) A$ is both bounded below and bounded above

- A bounded below $\nRightarrow A$ is bounded.
- A bounded above $\nRightarrow A$ is bounded.

Powers and roots:

$$2^2 = 4, \quad \sqrt{25} = 5 \dots$$

Let $x \in \mathbb{R}$, let $n \in \mathbb{Z}$

$$x^n := \begin{cases} \underbrace{x \dots x}_{n \text{ times}} \\ 1 \\ \frac{1}{x^{-n}} \end{cases}$$

$$n > 0$$

$$n = 0, x \neq 0$$

$$n < 0, x \neq 0$$

$$\lfloor \frac{1}{x^{-n}}$$

$$n < 0, x \neq 0$$

$$2^3 = 8, 2^{-3} = \frac{1}{2^3} = \frac{1}{8}, 3^2 = 9 \dots$$

. Roots:

. let $y \geq 0$, let $n \in \mathbb{N}$.

$$\sqrt[n]{y} = \sup \{ x \in \mathbb{R} : 0 \leq x^{2n} \leq y \} \geq 0$$

$$n=1, y=4, \sqrt[2]{4} = 2 \quad (n=2, \sqrt[3]{4} = \sqrt[3]{4})$$


. Properties:

$$(\sqrt[n]{y})^{2n} = y, \quad \forall y \geq 0.$$

$$\sqrt[n]{0} = 0$$


. Odd roots: we want to give a precise meaning, for example, to $\sqrt[3]{8} = 2$

$$\forall y \in \mathbb{R}, \text{ we set } |y| = \begin{cases} y & \text{if } y \geq 0 \\ -y & \text{if } y < 0 \end{cases}$$



$$\forall y \in \mathbb{R}, |y| \geq 0.$$

$$\forall y \in \mathbb{R}, \operatorname{sgn}(y) = \begin{cases} 1 & \text{if } y > 0 \\ 0 & \text{if } y = 0 \\ -1 & \text{if } y < 0 \end{cases}$$




$$\text{In particular, } \forall y \in \mathbb{R}, y = \operatorname{sgn}(y)|y|$$


$$\text{Take } y \in \mathbb{R}, n \in \mathbb{N}$$

$$\sqrt[2n+1]{y} := \operatorname{sgn}(y) \sup \{x \in \mathbb{R} : 0 \leq x^{2n+1} \leq |y|\}$$

$$\sqrt[3]{8} = 2, \quad \sqrt[3]{-8} = -2$$



even roots: always take +



odd roots: if $x > 0$ take +
if $x < 0$ take -

Remark :

$$(\sqrt[2n]{y})^{2n} = y \quad \forall y \geq 0$$

$$\cdot x \in \mathbb{R} \quad \sqrt[2n]{x^{2n}} = |x|.$$

$$x = -2, n = 1. \quad (-2)^2 = 4 \sim (\sqrt{-2})^2 = \sqrt{4} = 2 \neq -2$$

$$\cdot y \in \mathbb{R}, \quad \left(\sqrt[2n+1]{y} \right)^{2n+1} = y$$

$$\cdot x \in \mathbb{R} \quad \sqrt[2n+1]{x^{2n+1}} = x$$

Equation:


let $y \in \mathbb{R}, n \in \mathbb{N}$. We want to solve equation $x^{2n} = y$

Which are the solution?

$\cdot y < 0$: $x^2 = -1$. There is no solution.

$$\cdot y \geq 0: \quad x^{2n} = y \quad (==) \quad x = \pm \sqrt[2n]{y}$$

$$\left(-\sqrt[2n]{y} \right)^{2n} = (-1)^{2n} \left(\sqrt[2n]{y} \right)^{2n} = 1 \cdot y = y$$



we want to solve equation $y_0 = x^2$

$$y_0 > 0 \Rightarrow x = \pm \sqrt{y_0}$$

$$y_0 = 0 \Rightarrow x = 0$$

$$y_0 < 0 \Rightarrow \text{no intersection.}$$

$$x^2 = 4 \quad (\Rightarrow) \quad x = \pm 2$$

$$x^2 = 0 \quad (\Rightarrow) \quad x = 0$$

$$x^2 = -1 \quad \text{impossible.}$$

Odd exponents:


we want to solve $x^{2n+1} = y, \forall y \in \mathbb{R}.$

$$\Leftrightarrow \boxed{x = \sqrt[2n+1]{y}}$$

$\forall y \in \mathbb{R}$, equation $x^{2n+1} = y$ has a unique solution, which is given by $x = \sqrt[2n+1]{y}$

$n=1$, we want to solve $x^3 = y$

the solution is $x = \sqrt[3]{y}.$



$$x^3 = y_0 \quad (\Rightarrow) \quad \boxed{x = \sqrt[3]{y_0}}$$

Equations:

even powers:

$$x^2 = y$$

$$\begin{cases} y > 0 : x = \pm \sqrt{y} \text{ (2 solut)} \\ y = 0 : x = 0 \text{ (1 solut.)} \end{cases}$$

even powers: $x^2 = y$ $\left\{ \begin{array}{l} y = 0 : x = 0 \text{ (1 solut.)} \\ y < 0 : \text{no solution.} \end{array} \right.$

odd powers: $x^3 = y \Leftrightarrow x = \sqrt[3]{y}$ (unique sol)
 $\forall y \in \mathbb{R}.$