

DIPARTIMENTO DI MATEMATICA

Seminari di Matematica

Normalised solutions to a fractional Schrödinger equation in the strongly sublinear regime

Jacopo Schino

University of Warsaw, Poland

3 ottobre 2025, ore 15:00 in Aula VI

Abstract

Schrödinger-type equations model a lot of natural phenomena and their solutions have interesting and important properties: one of them is the conservation of mass, which gives rise to the search for normalisedsolutions. In this talk, I will explain a possible approach to solve

$$\begin{cases} (-\Delta)^s u + \mu u = g(u), \\ \int_{\mathbb{R}^N} u^2 dx = m, \\ (\mu, u) \in \mathbb{R} \times H^s(\mathbb{R}^N), \end{cases}$$

where $N \geq 2, \, 0 < s < 1,$ and m > 0 is prescribed, in cases that include the so-called strongly sublinear

regime: $\lim_{t\to 0} \frac{g(t)}{t} = -\infty. \tag{1}$ This makes a direct variational approach impossible because the energy functional is not well-defined in $H^s(\mathbb{R}^N)$. In the proposed approach, when m is sufficiently large, a family of approximating problems is considered so that the energy functional is of class C^1 and a corresponding family of solutions is obtained, which eventually converge to a solution to the original problem. When (1) holds, the previous result for a suitably translated problem is exploited to obtain a solution for any m>0. This is joint work with Marco Gallo (Catholic University of the Sacred Heart, Brescia, Italy).

Organizzato dalla Prof.ssa Silvia Cingolani e dal Prof. Gabriele Mancini nell'ambito delle attività del progetto PRIN PNRR 2022, P2022YFAJH - Linear and Nonlinear PDE's: New directions and Applications



PRIN PRINR 2022 P2022YFALH "Linear and Nonlinear PDE's: New directions and Applications"

Funded by the
European Use European PARO NR/DONALE DI RIPIESAR FRSLENKA (PRRIP) NewtitionerationEU

CIP History (CIP History Control of Component's Control of Component's Control of Con







